

NMA Survey of CO and HCN Emission from Nearby Active Galaxies

K. Kohno^{1*}, R. Kawabe¹, & B. Vila-Vilaró¹

¹Nobeyama Radio Observatory, Minamimaki, Minamisaku, Nagano, 384-1305, Japan

High resolution (a few arcseconds) observations of CO(1–0) and HCN(1–0) emission from nearby Seyfert galaxies have been conducted with the Nobeyama Millimeter Array. Based on the observed CO distribution and kinematics, we suggest that a small scale (a few 100 pc - a few kpc) distortion of the underlying potential seems to be necessary for Seyfert activity, although it is not a sufficient condition. We also find that Toomre’s Q values in the centers of Seyfert galaxies tend to be larger than unity, suggesting the circumnuclear molecular gas disks around Seyfert nuclei would be gravitationally stable. The HCN/CO integrated intensity ratios ($R_{\text{HCN/CO}}$) range over an order of magnitude, from 0.086 to 0.6. The Seyfert galaxies with high $R_{\text{HCN/CO}}$ may have an extended ($r \sim 100$ pc scale) envelope of obscuring material. The presence of kpc scale jet/outflows might be also related to the extremely high $R_{\text{HCN/CO}}$.

1. Differences between active and non-active galaxies

Molecular gas has been considered to be a promising fuel source which may be channeled into the central engines of active galaxies, and therefore the amount of molecular material in Seyfert galaxies is expected to be an important clue to understand the nature of active galactic nuclei. Surveys of CO emission from Seyfert galaxies with single-dish telescopes (*e.g.* Maiolino *et al.* 1997; Vila-Vilaró *et al.* 1998) find, however, no significant differences on the *total amount* of molecular gas between active and quiescent spirals. In order to deduce the conditions which are necessary for nuclear activity, we have conducted aperture synthesis observations of CO(1–0) and HCN(1–0) emission in nearby Seyfert galaxies. The proximity of the sample ($D < 25$ Mpc; Table 1) allows us to perform extensive studies of their molecular gas distribution, kinematics, and physical conditions at a few 100 pc scales.

2. Distributions and kinematics of molecular gas

Fig. 1 shows the integrated intensity maps and intensity-weighted mean velocity maps of the CO emission of Seyfert galaxies. The distributions of molecular gas show a wide variety, and it appears that there are no “typical” gas distributions either in type-1s or type-2s. It should be noted, however, that the observed features such as twin peaks (seen in NGC 4579, NGC 5033, and NGC 6951, for instance) have been often reported in the central regions of barred galaxies (*e.g.* Kenney *et al.* 1992), and all the observed gas distributions and kinematics seem to be responding to a non-axisymmetric potential (Kohno *et al.* 1999b).

It has been claimed that there exist Seyfert galaxies without bars (*e.g.* McLeod & Rieke 1995; Mulchaey & Regan 1997). However, very weak bars (a few % or less in density), which may be not easy to detect even in near-infrared bands, can drastically affect the behavior of the gas (Wada & Habe 1992). Moreover, critical elements of the fuel supply need not be evident at a large-scale in the host galaxy (McLeod & Rieke 1995). We therefore suggest that small scale (a few 100 pc - a few kpc) distortions of the underlying potential are necessary for the Seyfert activity, although it is *not* a sufficient condition (Sakamoto *et al.* 1998).

* e-mail: <kotaro@nro.nao.ac.jp>




Fig. 1: Distributions and kinematics of CO emission from the central regions of nearby Seyfert galaxies. The size of all CO maps is adjusted to the same linear scale in the figure. Contour levels are 2σ , 4σ , 6σ , $\dots$, for NGC 4051, NGC 4579, NGC 5033, and NGC 5194. Contour levels for NGC 3982 are 1.5σ , 3σ , 4.5σ , $\dots$, and 2σ , 5σ , 8σ , $\dots$, for NGC 6951. Noise levels (σ) are 3.3, 19, 6.7, 13, 14, and 15 K km s⁻¹ for NGC 3982, NGC 4051, NGC 4579, NGC 5033, NGC 5194, and NGC 6951, respectively. The cross in each map marks the location of Seyfert nucleus (6 cm radio continuum peak).

3. Gravitational stabilities of molecular gas

Self-gravity of molecular gas is another important issue because circumnuclear molecular gas can lose angular momentum to fall quickly into the nucleus if the gas is gravitationally unstable. The Toomre's Q parameters were therefore computed to examine the gravitational stabilities of the circumnuclear molecular gas disk in Seyfert galaxies (Fig. 2). We also used CO data in the literature to calculate Q values for non-Seyfert galaxies.

Fig. 2 shows that the molecular gas tends to be gravitationally *stable* near the Seyfert nuclei (Kohno *et al.* 1999b; Sakamoto *et al.* 1998). This result suggests that instabilities of molecular gas disk around the nuclei may not cause a further infall of gas in some Seyferts; if a circumnuclear gas disk becomes gravitationally unstable, it would result in a burst of star formation there, as in the case of pure starburst galaxies such as NGC 3504 and NGC 6946. Note that NGC 4051, which shows $Q < 1$, has an evidence for recent burst of star formation around the nucleus (Baum *et al.* 1993).

Perhaps our results might be also related to the fact that Seyfert nuclei preferentially reside in early type hosts; it is well known that early type galaxies have smaller gas mass - to - dynamical mass ratio (Young & Scoville 1992), and this means molecular gas in early type galaxies tends to




Fig. 2: Gravitational stabilities of circumnuclear molecular gas disks measured with the Toomre’s Q parameters. $Q \equiv \Sigma_{\text{crit}}/\Sigma_{\text{gas}}$, where Σ_{crit} is a threshold gas surface density for the onset of instabilities, and Σ_{gas} is an observed gas surface density. Σ_{crit} and Σ_{gas} are listed in Table 1 as well as Q values.

be gravitationally stable. Even in late type Seyferts such as NGC 5033, circumnuclear molecular gas disks may be similar to those of early type hosts in terms of gravitational stabilities.

4. Physical conditions of molecular gas

A striking enhancement of HCN(1–0) emission has been reported in the nuclear regions of some Seyfert galaxies such as NGC 1068 and NGC 5194 (Kohno *et al.* 1996 and references therein), implying extreme physical conditions of the molecular gas near active nuclei (*e.g.* Matsushita *et al.* 1998). Is an extremely enhanced HCN emission common or not in Seyfert galaxies?

The HCN/CO integrated intensity ratios ($R_{\text{HCN/CO}}$) in the observed Seyferts are listed in Table 1 together with $R_{\text{HCN/CO}}$ in non-Seyfert galaxies. The ratios range over an order of magnitude, from 0.086 to 0.6 among Seyfert galaxies. $R_{\text{HCN/CO}}$ values larger than 0.3 have never been reported in any non-Seyfert galaxies till date. We suggest that the observed $R_{\text{HCN/CO}}$ in Seyfert galaxies may be related to the size of *dense obscuring material*; NGC 1068 and NGC 5194 have a relatively extended ($r \sim 100$ pc scale) envelope of obscuring material, which consists of very dense interstellar medium and therefore shows extremely high $R_{\text{HCN/CO}}$. On the other hand, some Seyferts, NGC 6951 for example, may have only a small scale ($r < 10$ pc) obscuring torus; in this case, our observing beam would be too large to find any significant enhancement of $R_{\text{HCN/CO}}$ associated with the dense obscuring material.

Note that the “high $R_{\text{HCN/CO}}$ Seyferts”, i.e. NGC 1068, NGC 3079, and NGC 5194, possess kpc scale jet/outflows. Circumnuclear dense molecular materials may also play a role in the confinement of large scale jet/outflows (Scoville *et al.* 1998). See Kohno *et al.* (1999a, b) against the discussion on the chemical abnormality of molecular gas in Seyfert galaxies.

We are grateful to Dr. T. Helfer for sending her CO data of NGC 1068. K. K. was financially supported by the Japan Society for the Promotion of Science.

References

- Baum, S. A., O’Dea, C. P., Dallacassa, D., Bruyn, A. G., Pedlar, A. 1993, *ApJ* **419**, 553-572
Helfer, T., Blitz, L. 1995, *ApJ* **450**, 90-110
Jackson, J. M., Heyer, M. H., Paglione, T. A. D., Bolatto, A. D. 1996, *ApJ* **456**, L91-L94

Table 1: Observed properties of the molecular gas in the central regions of Seyferts and non-Seyferts

Name	Activity	Σ_{gas}	Σ_{crit}	Q	$R_{\text{HCN/CO}}$	Ref.
Seyfert Galaxies (type 1s and 2s)						
NGC 4051	Sy 1.2	1.8×10^3	1.6×10^3	0.90	...	(1)
NGC 5033	Sy 1.5	3.6×10^2	2.7×10^3	7.5	0.14U	(1)
NGC 1068	Sy 1.8	1.6×10^3	2.1×10^3	1.3	0.6	(2)
NGC 3079	Sy 2	2.5×10^3	...	...	0.3	(3), (1)
NGC 3982	Sy 1.9	2.1×10^2	9.1×10^2	4.2	0.20U	(1)
NGC 4579	Sy 1.9	5.2×10^2	...	...	...	(1)
NGC 5194	Sy 2	4.0×10^2	1.7×10^3	4.2	0.56	(4)
NGC 6951	Sy 2	8.2×10^2	1.7×10^3	2.1	0.086	(5)
Non-Seyfert Galaxies (Starbursts and Post-Starbursts)						
NGC 3504	Starburst	4.3×10^3	4.8×10^3	1.1	0.3	(6), (7)
NGC 6946	Starburst	5.5×10^3	3.4×10^3	0.61	0.12	(1)
NGC 4321	Hot-Spot	6.0×10^3	4.2×10^3	0.70	...	(8)
NGC 4736	Post-Starburst	7.6×10^2	3.6×10^3	4.8	0.044U	(7), (8)
NGC 7331	Post-Starburst	1.4×10^2 U	1.3×10^3	9.1L	...	(9)
Milky Way	Young Post-Starburst	1.1×10^3	3.6×10^3	3.3	0.08	(10), (11)

References. — (1) This work; (2) Helfer & Blitz 1995; (3) Sofue & Irwin 1992; (4) Kohno *et al.* 1996; (5) Kohno *et al.* 1999a; (6) Kenney *et al.* 1993; (7) Kohno *et al.* 1998; (8) Sakamoto *et al.* 1998; (9) Tosaki & Shioya 1997; (10) Oka *et al.* 1998; (11) Jackson *et al.* 1996.

Note. — (1) Galaxy name; (2) Type of activity; (3) Observed face-on gas surface density $\Sigma_{\text{gas}} \equiv 1.36\Sigma_{\text{H}_2}$ in $M_{\odot} \text{ pc}^{-2}$, adopting $X = 3.0 \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$. U means upper limit; (4) Critical gas surface density $\Sigma_{\text{crit}} \equiv \alpha(\sigma_v \kappa / \pi G)$ in $M_{\odot} \text{ pc}^{-2}$; (5) Toomre's Q value calculated by $\Sigma_{\text{crit}}/\Sigma_{\text{gas}}$. L means lower limit; (6) HCN to CO integrated intensity ratio in brightness temperature scale. U means upper limit. See also Kohno *et al.* (1999b); (7) References.

- Kenney, J. D. P., Wilson, C. D., Scoville, N. Z., Devereux, N. A., Young, J. S. 1992, *ApJ* **395**, L79-L82
Kenney, J. D. P., Carlstrom, J. E., Young, J. S. 1993, *ApJ* **418**, 687-708
Kohno, K., Kawabe, R., Tosaki, T., Okumura, S., K. 1996, *ApJ* **461**, L29-L32
Kohno, K., Kawabe, R., Sakamoto, K., Ishizuki, S., Vila-Vilaró, B. 1998, in “The Central Regions of Galaxy and Galaxies”, Sofue, Y. (ed.) 239-240
Kohno, K., Kawabe, R., Vila-Vilaró, B. 1999a, *ApJ* **511**, 157-177
Kohno, K., Kawabe, R., Ishizuki, S., Vila-Vilaró, B. 1999b, *Adv. Space Res.* **23**, 1011-1014
Maiolino, R., Ruiz, M., Rieke, G. H., Papadopoulos, P. 1997, *ApJ* **485**, 552-569
Matsushita, S., Kohno, K., Vila-Vilaró, B., Tosaki, T., Kawabe, R. 1998, *ApJ* **495**, 267-275
McLeod, K., K., Rieke, G. H. 1995, *ApJ* **441**, 96-112
Mulchaey, J. S., Regan, M. W. 1997, *ApJ* **482**, L135-L137
Oka, T., Hasegawa, T., Hayashi, M., Handa, T., Sakamoto, S. 1998, *ApJ* **493**, 730-761
Sakamoto, K., Okumura, S. K., Ishizuki, S., Scoville, N. Z. 1998, *ApJ*, submitted.
Scoville, N. Z., Yun, M. S., Armus, L., Ford, H. 1998, *ApJ* **493**, L63-L66
Sofue, Y., Irwin, J. A. 1992, *PASJ* **44**, 353-365
Tosaki, T., Shioya, Y. 1997, *ApJ* **484**, 664-671
Vila-Vilaró, B., Taniguchi, Y., Nakai, N. 1998, *AJ* **116**, 1553-1572
Wada, K., Habe, A. 1992, *MNRAS* **258**, 82-94
Young, J. S., Scoville, N. Z. 1992, *ARAA* **29**, 581-625